

**IMMEDIATE PUBLIC HEALTH CONCERNS
AND ACTIONS IN VOLCANIC ERUPTIONS:
LESSONS FROM THE MOUNT
ST. HELENS ERUPTIONS
May 18 - October 18, 1980**

Robert S. Bernstein, MD, PhD, Peter J. Baxter, MD, Henry Falk,
MD, MPH, Roy Ing, MD, MPH, Laurence Foster, MD, MPH,
and Floyd Frost, DrPH

Introduction

Before the eruptions of Mount St. Helens in 1980, little information was available in the clinical and environmental science literature on the health effects of volcanic eruptions.^{1,2} Some studies of volcanic hazards had been published in the literature of a variety of other disciplines (e.g., geology, sociology, psychology, and ecology)³⁻⁶; others were never published. Furthermore, despite the fact that Mount St. Helens was well known to earth scientists as the most active and dangerous volcano in the Cascade Range,⁷ there had been no explosive volcanic eruptions in the United States since 1914 (Mount Lassen), none at Mount St. Helens for more than 125 years, and no historical record of any fatalities due to explosive eruptions in the sparsely populated volcanic regions of the west coast of the United States and Canada.⁸

Thus, when Crandell, *et al*., predicted in *Science* in 1975 that Mount St. Helens, last active for a 27-year period from 1831 to 1857 "will erupt again—perhaps within the next few decades,"⁹ it was not too surprising that their report did not galvanize the local public health community.

After hazard prediction maps and disaster contingency recommendations were published in even more detail in a US Geological survey (USGS) Bulletin in 1978,⁹ several local and state agencies did review and update their contingency plans for flood-warning and evacuation, triage of mass casualties, and search and rescue operations. However, there was no apparent recognition of the desirability of coordinated predisaster consultation and planning for disaster response activities among government officials and technical expert

in the psychosocial, geological, and biomedical sciences together with law Enforcement officials and public health and safety officials.

As it turned out, the USGS volcanologists had been able to predict with considerable accuracy the types and geographic distribution of the principal safety hazards which ultimately resulted from the May 18 and subsequent eruptions of 1980. Unfortunately, they were unable to predict the precise times of onset or duration of any of the six major eruptions and they did not forecast the lateral direction and magnitude of the May 18 explosive blast and ashfall, or the earthquake and landslide which preceded in¹⁰

As a result, the May 18, 1980 eruption of Mount St. Helens caught volcanologists, government officials, and others largely unprepared for the nature and magnitude of its impact. Similarly, government- and university-based scientists were unprepared for the unprecedented and rapidly perishable opportunity to address various research needs concerning the evaluation and control of adverse health effects of explosive volcanism.

In this chapter, we provide a chronology of events leading to the May 18 eruption and a description of the impact of the ashfall on local communities which led to state and federal requests for epidemiologic assistance from the Centers for Disease Control (CDC). Following that, we describe the formation and actions of a coordinated state- and federal disaster response effort to develop accurate information about volcanic hazards and to recommend methods for prevention or control of adverse effects on safety and health. The purpose of this chapter is to provide a case study of the descriptive epidemiology of a major natural disaster. By studying the events preceding and resulting from the Mount St. Helens eruptions, high-risk communities in North America and elsewhere may be better able to address the issues concerning evaluation and control of adverse health effects of explosive volcanism and volcanic ash.

Chronology of Events and Response

Federal/State Responses

Premonitory activity began at Mount St. Helens on March 20, 1980 with an earthquake measured at 4.1 on the Richter scale. Earthquakes continued with increasing frequency until the first of a series of minor explosive eruptions occurred on March 27. Harmonic tremors, seismographic evidence of the movement of molten

rock beneath the mountain, began to occur on April 3 a few days after it was noted that an area on the north side of the summit had begun to bulge ominously by as much as 1.5 m m a day. By May 18, this deformation caused upward and outward displacement of up to 98 m per day and measured 1.6 by 1 km.

At 8:32 am on Sunday, May 18, a major earthquake, 5.0 on the Richter scale, caused the roof of the bulge to slide downhill, thereby permitting an estimated 50 billion liters of superheated water to expand into steam and escape in the form of an explosive blast directed toward the north. Fortuitously, hundreds of loggers who would have been authorized to be working within the area that was devastated were spared because the eruption occurred on a Sunday.¹⁰

Following the cataclysmic lateral blast of May 18, 1980, there were five additional major explosive eruptions (May 25, June 12, July 22, August 7, and October 16-18, 1980) and several non-explosive, "dome building" eruptions. Ashfalls of up to 70 mm or more in depth from four of the explosive eruptions were deposited by prevailing winds in sparsely populated areas to the north and east, while the ashfall of up to 10 mm from one blast and another of up to 5 mm were blown toward more populated centers to the west and southwest, respectively (Figure 1). The remaining 2.5 km of the volcano's cone provided an important "stack" effect in dissipating volcanic gases¹¹ (principally SO₂) which decreased gradually from the initial daily rates of more than 1,000 tons.^{12,13} Within hours of the eruption, it was clear that mudflows were likely to cause flooding and blockage of waterborne transport downstream from the volcano.^{10,14 (12)*} In communities affected by heavy ashfalls, the mineral dust was inadvertently washed into water treatment plants where it occasionally overburdened filters and damaged machinery.^{14(4 5.9. 10.12)}

As one would expect, there were widespread and significantly adverse socioeconomic effects due to the mudflows and resultant flooding in local communities.¹⁵⁻²¹ However, the question most often posed to physicians and public health officials was, "Is the volcanic ash harmful to health?"^{22 23}

On May 19, 1980, State Health Department officials in Idaho telephoned the Center for Environmental Health (CEH) at CDC requesting information on the health effects of volcanic ash because of widespread public concern over the recent ashfalls. As a result, on




FIGURE 1.—Ashfall after just three major eruptions of Mount St. Helens and locations of Washington and Oregon hospitals under Centers for Disease Control surveillance. Ashfall paths: May 18, 1980, northeasterly affecting Yakima, Ellensburg, and Spokane, Washington; May 25, northwesterly affecting Chehalis and Centralia, Washington; June 12, southwesterly affecting Vancouver, Washington and Portland, Oregon.

SOURCE: US Geological Survey professional paper #1250. Reprinted with permission from the *Journal of the American Medical Association*. *

May 20, 1980, environmental health scientists from CEH and the National Institute for Occupational Safety and Health (NIOSH), CDC, met with officials from the Federal Emergency Management Agency (FEMA), Environmental Protection Agency (EPA), USGS, United States Forest Service (USFS), National Oceanographic and Atmospheric Administration (NOAA), and the National Weather Service, as well as other concerned federal agencies, to assess the apparent disaster in the north west. It was subsequently learned that from May 18 through May 21 the 24-hour average concentrations of total suspended particulates (TSP), monitored by EPA's roof-top air monitoring stations in Yakima and Spokane, Washington, and Coeur d'Alene, Idaho, persistently exceeded the "warning levels" ($625 \mu\text{g}/\text{m}^3$) of EPA's National Ambient Air Quality (NAAQ) Standards for TSP derived from fossil fuel pollutants (see Table 1). 14(3,4,9),24, 25

On May 21, 1980, at the request of the State of Washington, the President declared the State a disaster area. The State Health Officer

had already requested epidemiologic assistance from CEH and NIOSH, CDC. As a result, between May 21 and September 13, 1980, more 30 CDC professional—including medical epidemiologists, industrial hygienists, engineers, statisticians, nurses, technicians, and interviewer—conducted field studies and provided technical consultation to state and federal agencies in Washington and Oregon, while others were engaged in laboratory studies and support activities at CDC facilities. 26–30

Community Response to Public Health Concerns

In Yakima, a town of more than 50,000 residents in an irrigated desert valley approximately 135 km northeast of Mount St. Helens, 24-hour average concentrations of TSP during the week after the May 18 eruption persistently exceeded the EPA "significant harm" ($1000 \mu\text{g}/\text{m}^3$) NAAQ Standards designed for monitoring and control of fossil fuel air pollution (see Table 1). The air quality problem was exacerbated by continual resuspension of sedimented volcanic ash by wind or by automobile traffic.

Yakima residents and public health officials were caught unprepared for the eruption, the ashfall, and the total darkness which engulfed the town by late morning. They were alarmed by the irritant effects of the ash on the eyes and respiratory tract and worried

--	--	--	--	--

TABLE 1—Results of Environmental Protection Agency (EPA) Air Monitoring for Total Suspended Particulates (TSP) Before and After the May 18 Eruption, 24-Hour Average Concentrations ($\mu\text{g}/\text{m}^3$), Yakima, Washington, 1980*

Before May 18	On May 18	May 19–May 25	On May 26	May 27–June 11
≤ 50	33,400	5,800–13,000	250	50–250

*In the early morning of May 26, a prolonged and heavy rainfall occurred in the town of Yakima. Ashfalls from the major eruptions of May 25 and June 13, 1980, were deposited to the west and southwest of Mount St. Helens, respectively (i.e., in the opposite direction from Yakima).¹⁴ (3–4–9) The EPA Action Levels for 24-hour average concentrations of TSP, derived from the combustion of fossil fuel pollutants, are: Alert, $375 \mu\text{g}/\text{m}^3$; Warning, $625 \mu\text{g}/\text{m}^3$; Emergency, $875 \mu\text{g}/\text{m}^3$; and Significant Harm, $1000 \mu\text{g}/\text{m}^3$.²⁵

--	--	--	--	--	--

about the possible adverse effects on water quality, agricultural products, farm animals, and machines. 10,16,17 Furthermore, the ash, which smelled of sulfur, severely limited visibility and led to motor vehicle accidents. 14(5,15).

Every able-bodied person was mobilized to clean up the ash, using wetting methods to reduce the resuspension of dusts and improvising makeshift masks of every sort—including wet or dry cloths, surgical masks, disposable industrial masks, and elaborate filtered- and supplied-air respirators—to reduce the concentration of inhaled particles.¹⁰

Throughout the week from May 18–26, volcanic ash fell in Washington and Idaho on a total area containing more than a million people, many of whom sought advice from physicians and public health officials. The most commonly expressed concerns were about the short and long-term hazards of inhalation of ash, ingestion of ash-contaminated food and water, and appropriate and effective methods of protection.

A team of CDC, state, and university-based environmental health scientists visited Yakima on May 25-26, guided to the town by local officials through a blinding dust storm. Discussions with emergency room physicians and preliminary reviews of hospital records confirmed news reports and passive surveillance data concerning apparent increases in respiratory morbidity. These findings suggested the need for detailed epidemiologic surveillance and epidemic field investigations (as outlined in Table 2), 31 On May 26, a steady rainfall in the Yakima area brought relief by purging the air of ash and aiding cleanup operations; however, concerns about the effects of volcanic ash on public health remained. Accordingly, refinement of the epidemiologic surveillance system and organization of more detailed studies were rapidly accomplished as described below.

State/Federal Disaster Response

It was immediately apparent after the May 18, 1980 eruption that a coordinated effort among many different agencies was necessary in order to protect the public's health and safety. 32–38 State public health officials were presented with important problems of identifying, evaluating, and controlling a unique new set of acute health and safety hazards. Simultaneously, there was the opportunity to carry out applied and basic science research to identify and

TABLE 2—Outline for Epidemic Field Investigations of Suspected Outbreaks

1.	Confirm Epidemic Occurrence and/or Distribution Cases <ul style="list-style-type: none"> • Review medical records and available epidemiologic surveillance data • Discuss with federal and state or local public health investigators
2.	Verify Diagnoses of Index Cases <ul style="list-style-type: none"> • Examine patients and medical and laboratory reports, possible • Discuss diagnostic criteria with reporting health professionals
3.	Develop Standardized Case Definition(s) <ul style="list-style-type: none"> • Define criteria for definite, probable, and suspect cases • Include clinical, laboratory, and epidemiologic criteria • Characterize potential etiologic exposures by quantitative measures
4.	Seek and Investigate Additional Cases (Numerator Data) <ul style="list-style-type: none"> • Ask index cases about other affected individuals • Establish a "hotline" telephone number, using public media • Review records of hospitals and physicians in expanding circle • Conduct surveys of affected communities or workplaces
5.	Develop a Rough Case Count and Line-Listing of Descriptive Data
6.	Orient the Descriptive Data on Cases by Person, Place, and Time <ul style="list-style-type: none"> • Prepare an epidemic curve and other graphic representations of data
7.	Determine Who is at Risk of Becoming Affected (Denominator Data)
8.	Develop an Etiologic Hypothesis and Appropriate Statistical Methods for Testing its Validity <ul style="list-style-type: none"> • Consider opportunities for prevention and/or control • Explain source and mode of transmission of putative agent(s)
9.	Analyze Descriptive Data: Compare the Hypothesis with Established Facts <ul style="list-style-type: none"> • Identify possible host-related and environmental risk factors • Identify possible sources, reservoirs, vectors, and agents
10.	Implement Interim Control Measures and Refine Epidemiologic Surveillance
11.	Analyze Surveillance Data and Conduct Additional Studies as Necessary <ul style="list-style-type: none"> • Refine case definition(s) and case-finding procedures • Determine the need for case-control or longitudinal studies • Identify a suitable reference or comparison population
12.	Prepare and Disseminate Timely Reports of Findings and Recommendations
13.	Evaluate the Effectiveness of Control and/or Prevention Measures
14.	Develop and Implement Refined Control and/or Prevention Measures

TABLE 3—Concerns and Actions of Public Health Agencies Following the May 18, 1980 Eruption of Mount St. Helens

State Health Departments' Concerns About Needed Actions	Federal and State Public Health Agencies' Coordinated Responses to These Concerns
1. Assessment of potential health impact and recommendations about prevention and control	<p>Rapid establishment of community-based active surveillance system for cause-specific increases in emergency room visits and hospital admissions.</p> <p>Epidemic field investigations of suspected eruption-related outbreaks.</p> <p>Assessment of factors associated with death or survival with and without injuries for persons within 40 km of Mount St. Helens' crater, May 18, 1980.</p> <p>Initiation of cross-sectional, case-control, and longitudinal studies of adverse respiratory effects among high-risk (hypersusceptible or heavily exposed) groups.</p>
2. Restricted areas—designation and enforcement near Mount St. Helens, for the general public and for logging operations.	Laboratory studies of the toxicity of volcanic ash by <i>in vitro</i> and <i>in vivo</i> methods.
3. Public warning system—premonitory events, eruptions, or adverse secondary events such as floods, mudflows, or ashfalls requiring evacuation or other primary preventive measures.	<p>Continuous reassessment of restricted area boundaries, nature of restrictions, and risk factors for adverse effects.</p> <p>Review of literature concerning volcanic hazards effects on public health and safety.</p> <p>Dissemination of accurate public health information, including investigation and control of rumors, in coordination with officials from local, state and federal agencies, and private institutions.</p>
4. Air monitoring—nature and concentration of respirable and/or total suspended particulates in ashfall areas, determination of sampling methods and locations, and interpretation of data	Systematic collection and analysis of sedimented and airborne volcanic ash and estimation of human exposures in environmental and occupational settings.
5. Protection of drinking water systems located downstream from sewage treatment plants operating in a bypass mode due to ash sediment.	Systematic collection and analysis of waterborne concentrations of ash leachates and evaluation of the chemical and biological quality of drinking water supplies.

control the potential adverse chronic health effects associated with the eruption. Furthermore, it was likely that the explosive activity would continue intermittently, perhaps for year, 7-9 and valid and reliable data were needed to provide timely advice to clinicians and the public on the nature and prevention of hazardous exposures. In view of the limitations on human and technical resources available at the local and state levels, requests for federal epidemiological assistance were met with appropriate responses (Table 3).

The US Public Health Services is one of the leading Federal agencies to respond to emergency preventive health needs resulting from natural and man-made disasters. In this capacity, CDC provides epidemiological and technical health-related assistance to other federal agencies and to state health departments upon request. By providing assistance of this type, the CDC works to meet the goals of the Surgeon General for promoting health and preventing disease. 39

Hospital-Based Surveillance Network

The main hospitals in affected areas of eastern, central, and western Washington State were coordinated in a systematic assessment of emergency room (ER) visits and hospital admission for respiratory and other health and safety problems. This was done initially by active telephone surveillance of daily totals for selected cause-specific ER visits and admissions. Later, hospitals permitted CDC and state health department professionals to have access to hospital records for epidemic field investigations and detailed reviews of those patients attending the surveillance hospitals after April 1, 1980 (to provide some baseline, pre-eruption data on ER visits and admissions). 14.26

As an example, a list of diagnostic labels and the number of patient visits to one of the ash-affected ERs is shown in Table 4. These surveillance data were the first to indicate potential eruption-related problems requiring epidemic field investigations by a team of state and federal public health professionals. Since these are numerator data and the numbers are generally small, it is important not to overinterpret any apparent increases or decreases in specific diagnostic categories (e.g., visits for ear problems and foreign bodies in the eyes) without first confirming the surveillance findings in other communities with comparable exposures or carrying out more detailed field studies. In Table 2, we have provided an outline of the

TABLE 4—Reasons for Emergency Room Visits in Moses Lake, Washington, by Week, May 4–31, 1980

Reason for Visit	5/4–10	5/11–17	5/18–24	5/25–31	% Change
Accidents/injuries, total	61	63	43	45	–29
Motor vehicle accidents	2	4	11	5	+117
Falls	1	3	6	3	+125
Other injuries	58	56	26	37	–45
Respiratory, total	8	9	27	21	+182
URI	1	1	6	9	+650
Pneumonia/influenza	0	1	1	1	+100
Asthma	2	1	8	5	+333
Bronchitis	2	2	3	4	+75
COPD/emphysema	2	2	3	2	+25
Other pulmonary	1	2	6	0	+71
Ear (otitis)	1	0	6	1	+600
Eye (foreign bodies)	9	8	9	6	–12
Psychiatric	3	1	3	0	–25
All others	125	140	212	152	+39
Overall Total	207	221	300	225	+23

*Per Cent increase or decrease in ER visits for May 18–31, 1980, compared with visits for May 4–17, 1980. The first volcanic eruption with ashfall in Moses Lake (30- to 70-mm depth) occurred on May 18, 1980.¹⁴⁵⁾

procedures to be followed in such epidemic field investigations; these procedures are described in more detail elsewhere.^{31,40}

The surveillance network proved valuable in several ways. It made possible the confirmation of physicians' impressions that there was no convincing evidence of excessive ash-related *mortality* in affected communities due to cardiopulmonary problems, traffic accidents, flooding, or other causes. However, transient increases in *morbidity* due to motor vehicle accidents and falls were seen in some communities affected by heavy ashfalls.

Undue anxiety did not appear to occur in excess nor did severe eye injuries, although ashfall-related visits for mild anxiety and eye injuries did occur.^{14,41} The ability to detect subtle mental health effects was limited because data on such effects were collected in hospital ERs and not in community mental health facilities or by private mental health practitioners. Chapter 9 presents the results of a detailed community-based study of mental health effects.

No increases in communicable diseases were detectable, although delayed effects of the eruptions on the local ecology were subsequently associated with an outbreak of giardiasis.⁴² It is not clear whether this represented a true outbreak or a pseudoepidemic due to the increased intensity and quality of surveillance activities.

Transient increases in ER visits for acute effects on the respiratory tract (primarily asthma and acute bronchitis) were consistently observed following ashfall. These increases were significantly increased in affected communities of all sizes, not only in the smaller communities where doctors' offices were temporarily closed; ERs were the predominant resources for provision of medical services (Figures 2 and 3). 14.23.26

The apparent absence of any excess of respiratory mortality due to the five-day period of extremely high levels of TSP may have been due to several factors: 1) the precautions taken by susceptible members of the community as a result of advisories from physicians and the Washington Lung Association to avoid exposure to the ash; 2) the relatively less severe adverse effects which may be experienced from exposure to high levels of TSP alone versus the combined effects of exposure to TSP, complex organic chemicals, and metal fumes in fossil fuel emissions; and 3) the limited sensitivity to detect such an excess by hospital-based surveillance data alone. 43.50)

Of course, a surveillance system is never completely accurate. There were several limitations to the surveillance system established in the week following the May 18, 1980 eruption of Mount St. Helens. For example, no pre-eruption plan had been prepared for standardizing data collection, for defining ash-related adverse respiratory effects,⁵¹ or for routine recording of such information as occupation, smoking, and intensity, frequency, and duration of exposure to ash. In an ongoing surveillance system, it may be possible to make refinements which will provide more information about specific health and safety problems in relation to well characterized exposures to volcanic hazards.

However, it is the distribution and trends in reporting of diseases or injuries that, upon routine analysis of surveillance data, are most important. 35.52 Even if the surveillance system captures only a small proportion of the incident cases, as long as the system and the population at risk do not change substantially, it should provide reliable data about trends in the incidence of specific safety and health problems.

Public health officials can monitor these trends to determine the need for epidemic field investigations and more detailed epidemiologic studies to evaluate the effectiveness of various control measures. In the wake of the eruptions of Mount St. Helens, several types of

epidemiologic studies were carried out to develop accurate information about risk factors for adverse effects on safety and health in volcanic eruptions. These are described below.

Assessment of Death or Survival Factors

Relatively low mortality—35 known deaths, and 23 missing persons presumed dead—has been associated to date with the initial and five subsequent explosive eruptions. Investigators from CDC reviewed the autopsy findings, investigated the circumstance surrounding the deaths, and interviewed 100 survivors to determine factors related to death or survival. Results are summarized in Table 5**

All known deaths were directly attributable to the blast, pyroclastic flow, mudflows, and ashfall of the May 18 eruption. All but two of the 58 dead and missing had been in the areas of blast, mudflow, tree blowdown, and tree damage which occurred within a wedge-shaped sector extending 10–19 km from the crater in a northerly direction (Figure 4 and Table 5).




FIGURE 2—Daily temperatures (maximum and minimum), average wind velocities, 24-hour average concentrations of total suspended particulates (TSP) ($\mu\text{g}/\text{m}^3$), and number of emergency room visits due to asthma and bronchitis at the two major hospitals in Yakima, May 4 to June 2, 1980. The Environmental Protection Agency's Emergency Action Level for potential health problems due to increased air pollution (derived from the combustion of fossil fuels) is a 24-hour mean concentration of 875 $\mu\text{g}/\text{m}^3$ of TSP.
Reprinted with permission from *Archives of Environmental Health*¹⁴

Only 17 known survivors had been within the damaged areas, and all of them had been at the edge of the blowdown area or even farther away from the crater in the area of damaged timber. Thus, in the 19 x 40 km wedge-shaped area of blast, mudflows, and tree blowdown and damage, the overall mortality rate was 48/65 or about 74 per cent (approximate 95 per cent confidence limits = 62 per cent to 83 per cent). Twelve (71 per cent) of the survivors in the damaged areas suffered from serious injuries—six with second degree burns, and six with adverse respiratory effects due to ash inhalation.

Despite the establishment of “red” (no public access) and “blue” (daylight entry with special permit) danger zones on the basis of the best available information prior to the May 18 eruption, all of the victims were recovered from outside (or were authorized to be within) the danger zones (Figure 4). 14(10,19),53,54 4 Beyond the tree destruction zone to the north, two deaths had occurred due to ash inhalation, and 37 survivors were identified, giving an overall mortality rate of 2/37 or about 5.4 per cent (approximate 95 per cent confidence limits = 1.5 per cent to 18 per cent). All of these survivors—as well as 48 additional survivors located south of the volcano out of the path of the lateral blast, pyroclastic and mud flows, or heavy ashfall—escaped without serious injury.



Among the 25 victims who were autopsied, suffocation by airway obstruction with ash was the cause of death in 17 and contributory in two others with thermal injuries (76 per cent); thermal injuries alone accounted for three victims (12 per cent) died of trauma, primarily head injuries. 53,54

Although other interpretations are possible, the investigators reached the following conclusions: 14(11,19),53,54,55(3)

- o Mortality and severe injuries were highest in the zones of greatest physical damage, including areas to which commercial and public access were permitted under specific circumstances.
- o Within each damage zone, mortality and injuries were increased with increasing proximity to the mountain.
- o Enclosed vehicles provided some degree of protection against exposures to the blast, heat, projectiles, and ash.
- o Within comparable damage areas, survivors with least injuries appeared to be those who: were familiar with the area and had planned escape routes; were close to their vehicles and departed promptly;

and/or improvised protective measures against burns and ash inhalation.

o Medical personnel caring for evacuees and survivors should be prepared to adequately treat infected burn wounds and respiratory tract injuries.






FIGURE 3—Weekly total numbers (April to June 1980) of main airways-related logbook diagnoses for emergency room visits at two hospitals in Yakima (May 18 eruption, 4-mm ashfall) (top left), two hospitals in Chehalis and Centralia (May 25 eruption, 3-mm ashfall) (top right), and three hospitals in Portland (May 25, trace of ash, and June 12, 3-mm ashfall) (bottom). Reprinted with permission from the *Journal of the American Medical Association*²⁵

TABLE 5—Deaths, Missing Persons, Survivors, and Mortality by Damage Zones, May 18, 1980, Eruption of Mount St. Helens

Damage Zones	Deaths	Missing (Presumed Dead)	Survivors	Total	% Mortality
Blast, Mudslides	1	10	2	13	85
Downed Timber	30	5	7	42	83
Damaged Timber	1	1	8	10	20
Intact Timber	2	0	35*	37*	5 (2*)
Unknown	1	7	—	8	—
Total	35	23	52*	110*	53 (37*)

*Excluding 48 survivors located on the south side of the mountain.

+ In parentheses is the % mortality including in the denominator the 48 survivors on the south side of the mountain (no eruption-related deaths occurred south of the mountain).

Collection, Analysis of Volcanic Ash

While public safety officials were engaged in finding and recovering survivors and victims in the devastated "blowdown" area north of Mount St. Helens, environmental health scientists were investigating the nature and potential toxicity of the volcanic ash. 22,23 Flooding and other safety hazards of an eruption of Mount St. Helens had been anticipated and perceived in advance as real risks by local residents and health officials, but extensive ashfalls had not been widely anticipated or perceived as major risks. 15,18-20, 56,57 As a result of pre-eruption planning and coordination between the National Weather Service and the state and local safety officials, residents of communities threatened by mudflows and flooding were evacuated within hours of the May 18 eruption. 57 However, the lack of anticipation of heavy ashfalls left safety and health officials with no pre-disaster plans for respiratory protection, hospital-based surveillance of cardiopulmonary problems, coping behavior by individuals with preexisting chronic respiratory disease, or ash clean-up and disposal. 15,18,20

Dry sedimented samples of the May 18 volcanic ashfalls were collected from Spokane and Yakima by state laboratory personnel and NIOSH industrial hygienists, taking care not to include soil or road dusts. Preliminary NIOSH analyses of four samples were reported on May 30 and June 3, 1980 14,55 as follows (chapters 5 and 6 present more detailed information):

